

AMENDMENTS TO THE CLAIMS:

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (currently amended) ~~Process~~ A process for detecting and/or quantifying non-covalent interactions between a target protein and one of its ligands, comprising:

- preparing cells or cell fragments containing a ~~[[DNA]]~~ nucleic acid sequence ~~comprising a gene~~ expressing a fluorescent protein, fused with a ~~[[gene]]~~ nucleic acid for the target protein, the fusion between the ~~[[gene]]~~ nucleic acid for the fluorescent protein and the gene for the target protein being ~~[[such]]~~ so that the properties of the target protein are not modified by the presence of the fluorescent protein, wherein the interaction between the target protein, and the ligand is not modified, and wherein a response transduction function is not modified, the fluorescent protein ~~being chosen~~ is selected from ~~[[the]]~~ fluorescent proteins obtained or derived from autofluorescent proteins of cnidarians, the molar extinction coefficient of which is greater than about $14,000 \text{ M}^{-1}\text{cm}^{-1}$ and the quantic fluorescence yield is greater than about 0.38, this protein being selected from the group consisting of green fluorescent protein (GFP), ~~variants of GFP that conserve the fluorescence property, and fragments of GFP that conserve the~~

~~fluorescence property,~~ placing said cells or said cell fragments in contact with a ligand for said target protein, said ligand labeled with a ~~label comprising a~~ molecule capable of absorbing the light emitted by the fluorescent protein, or a fluorescent substance, the fluorescent protein being the fluorescence energy donor and the label being the fluorescence energy acceptor, or the fluorescent protein being the fluorescence energy acceptor and the label being a fluorescent substance which is a fluorescence energy donor, and irradiating said cells or said cell fragments at a wavelength which makes it possible either to excite the fluorescent protein or to excite the fluorescent substance, wherein the steps of placing in contact and irradiating are carried out either simultaneously or one after the other, or said cells or said cell fragments are placed in contact with a ligand for said protein, said ligand labeled with a label, the cells or the ligand having been irradiated before being placed in contact, and

- wherein a reduction in the amplitude of the donor's emission and/or emission signal characteristic of the acceptor's emission is ~~detected~~ measured and measuring the fluorescence energy transfer ~~when quantifying~~ to detect and/or quantify the non-covalent interactions.

2. (currently amended) ~~Process~~ The process according to Claim 1, wherein the protein whose protein-ligand interaction is selected from the group consisting of:

- membrane-bound proteins coupled to the G protein,
- growth factor receptors which are structurally linked to the insulin receptor,
- ion channel-receptors, and
- intracellular nuclear receptors which are structurally linked to the steroid receptor.

3. (currently amended) ~~Process~~ The process according to Claim 1, wherein the fluorescent protein is [[EGFP]] Enhanced Green Fluorescent Protein (EGFP) and the labeled substance is Bodipy and in which either the reduction in the emissions amplitude of EGFP or an emission signal of Bodipy resulting from an energy transfer is detected, the irradiation wavelength corresponding to the excitation wavelength of EGFP.

4. (currently amended) ~~Process~~ The process according to Claim 1, wherein the fluorescent protein is [[EGFP]] Enhanced Green Fluorescent Protein (EGFP) and the labeled substance is a coumarin, and wherein either the diminution of amplitude of coumarin or an emission signal EGFP resulting from an energy transfer is detected, said irradiation wavelength corresponding to the excitation wavelength of coumarin.

5. (currently amended) ~~Process~~ The process according to Claim 1, wherein the fluorescent protein is fused on the N-

terminal side and the target protein is fused on the C-terminal side.

6. (currently amended) ~~Process~~ The process according to Claim 1, wherein the fluorescent protein is fused on the C-terminal side and the target protein is fused on the N-terminal side.

7. (currently amended) ~~Process~~ The process according to Claim 1, wherein the fluorescent protein is inserted into a receptor target protein wherein said receptor is coupled to the G protein, this insertion taking place in the first or third intracellular loop of the receptor, with the proviso that the insertion does not destroy either the properties of the receptor or the fluorescence of the fluorescent protein.

8. (currently amended) ~~Process~~ The process according to Claim 1, wherein the cells are mammalian cells, ~~in particular HEK 293 cells~~ which are adherent or in suspension selected from the group consisting CHO cells, COS cells, lymphocytic lines, fibroblasts, yeast cells *pichia pastoris*, *saccharomyces cerevisia*, *saccharomyces kluyveri*, *Hansenula polymorpha*, and insect cells ~~infected with a virus and fungi~~.

9. (currently amended) ~~Process~~ The process according to Claim 1, wherein a signal can be detected, ~~in a conventional fluorimetry device or in a rapid-mixing device equipped with a system for detecting fluorescence,~~ after mixing the donor and the acceptor, and can be abolished by the addition of a non-

fluorescent substance of similar pharmacological specificity, and wherein a signal/noise ratio is greater than about 2.

10. (currently amended) ~~Process~~ A process for detecting and/or quantifying non-covalent interactions between a target protein ~~consisting of~~ comprising a receptor coupled to G proteins and a G protein, ~~in order~~ to identify [[the]] molecules which are biologically active with respect to the receptor, and which are capable of forming a reversible non-covalent interaction with said receptor, wherein:

- cells or fragments of cells which express a DNA sequence ~~comprising the gene coding~~ encoding for a fluorescent protein and fused with ~~the gene for the~~ nucleic acid expressing a receptor coupled to the G proteins are prepared, the fusion between the ~~gene coding~~ nucleic acid encoding for the fluorescent protein and the [[gene]] nucleic acid encoding for said receptor being that the properties of the receptor are not modified by the presence of the fluorescent protein, namely:

- * the interaction between the receptor and the G protein is not modified,

- * the interaction between the receptor and the biologically active molecule is not modified,

- * a response transduction function is not modified, the fluorescent protein ~~being chosen~~ is selected from the fluorescent proteins obtained or derived from autofluorescent proteins of cnidarians, the molecular

extinction coefficient of which is greater than about 14,000 $M^{-1}cm^{-1}$ and the quantic fluorescence yield of which is greater than about 0.38, this protein being ~~chosen in particular from a:~~

- green fluorescent protein (GFP), ~~or~~

~~variants derived from GFP by addition, deletion or substitution of one or more amino acids, with the proviso that these variants conserve the fluorescence property,~~

~~or fragments of GFP, or fragments of the above-mentioned variants, with the proviso that these fragments conserve the fluorescence property,~~

wherein the G protein ~~being~~ is labeled with a label ~~consisting comprising:~~

- either ~~[[of]]~~ a molecule which is capable of absorbing the light emitted by the fluorescent protein,

- or ~~[[of]]~~ a fluorescent substance,

~~[[-]]~~ wherein the fluorescent protein and ~~the above-mentioned said~~ label being such that they transfer energy from one to the other, ~~it being possible for wherein~~ the fluorescent protein ~~to be~~ is an energy donor or ~~it being possible for the above-mentioned said label to be~~ is an energy donor,

detecting the interaction between the receptor labeled with the fluorescent protein and the G protein labeled with said label by fluorescence energy transfer and measuring the

fluorescence energy transfer when quantifying the non-covalent interactions.

11. (currently amended) ~~Process~~ A process for identifying and/or quantifying interactions between a receptor and a non-fluorescent molecule which is biologically active with respect to ~~[[the]]~~ said receptor, which ~~[[are]]~~ is capable of forming a reversible non-covalent interaction with said receptor, by implementing the process ~~defined~~ according to claim 10, in which ~~[[are]]~~ a biologically active non-fluorescent molecule is added to cells, or cell fragments, which express the DNA coding for the receptor labeled with the fluorescent protein and for the G protein labeled with the label, characterized in that:

- an agonist and biologically active non-fluorescent molecule triggers a signal transduction detected by variation in the energy transfer between the receptor labeled with the fluorescent protein and the G protein labeled with the label;

- an antagonistic biologically active non-fluorescent molecule inhibits the signal transduction brought about by an agonist and detected by variation in the transfer of fluorescence energy between the receptor labeled with the fluorescent protein and the G protein labeled with the label.

12. (currently amended) ~~[[Kit]]~~ A kit or equipment for detecting and/or quantifying non-covalent interactions between a target protein labeled with a fluorescent protein and one of its ligands labeled with a label consisting:

- a molecule which is capable of absorbing the light emitted by the fluorescent protein,

- or a fluorescent substance,

this fluorescent protein ~~being chosen~~ is selected from [[the]] fluorescent proteins obtained or derived from autofluorescent proteins of cnidarians, the molecular extinction coefficient of which is greater than about $14,000 \text{ M}^{-1}\text{cm}^{-1}$ and the quantic fluorescence yield of which is greater than about 0.38, this protein being further chosen from green fluorescent protein (GFP), ~~variants of GFP that conserve the fluorescence property, and fragments of GFP and said variants,~~ that conserve the fluorescence property and its ligand labeled with a fluorescent substance, the said kit comprising:

- the target protein fused with a fluorescent protein or a stable cell line which is capable of expressing the protein fused with a fluorescent protein or a plasmid containing the nucleic acid sequence coding for [[the]] said targets protein fused with a fluorescent protein as defined above,

- the ligand labeled with ~~the above-mentioned~~ said label,

- buffers and media required for the energy transfer between said protein and said ligand.

13. (currently amended) [[Kit]] A kit or equipment for detecting and quantifying non-covalent interactions between a target protein labeled with a first fluorescent protein and one of

its ligands labeled with a fluorescent substance corresponding to a second fluorescent protein, said first fluorescent protein being chosen from the fluorescent protein enhanced yellow fluorescent protein (EYFP) or enhanced green fluorescent protein (EGFP) and the ligand being labeled with said second fluorescent protein enhanced cyan fluorescent protein (ECFP), or said first fluorescent protein being ECFP and the ligand being labeled with said second fluorescent protein EYFP or EGFP, [[the]] said kit comprising:

- a plasmid containing a nucleic acid sequence coding for the target protein fused with a fluorescent protein, and a plasmid containing a nucleic acid sequence coding for the ligand fused with a second fluorescent protein, or a ligand fused with a second fluorescent protein, obtained via a recombinant route and purified, or

- a stable cell line which is capable of expressing the target protein fused with first fluorescent protein, and a stable cell line which is capable of expressing the ligand fused with a second fluorescent protein or a ligand fused with a second fluorescent protein, obtained via a recombinant route and purified, and

- buffers and media required for an energy transfer between said protein and the said ligand.

14. (currently amended) ~~[[Kit]]~~ A kit or equipment for detecting and quantifying non-covalent interactions between a target protein consisting of a receptor coupled to the G protein labeled with a first fluorescent protein and ~~[[the]]~~ G protein labeled with a fluorescent substance corresponding to a second fluorescent protein, the first fluorescent protein being chosen from the fluorescent protein EYFP or EGFP and the G protein being labeled with the second fluorescent protein ECFP or the first fluorescent protein being ECFP and the G protein being labeled with the second fluorescent protein EYFP or EGFP, ~~[[the]]~~ said kit comprising:

- a plasmid containing a nucleic acid sequence coding for the receptor fused with a first fluorescent protein, and a plasmid containing a nucleic acid sequence coding for the G protein fused with a second fluorescent protein, or the G protein fused with a second fluorescent protein, obtained via a recombinant route and purified; or

- a stable cell line which is capable of expressing the receptor fused with a first fluorescent protein, and a stable cell line which is capable of expressing the G protein fused with a second fluorescent protein, or the G protein fused with a second fluorescent protein second , obtained via a recombinant route and purified; and

- buffers and media required for the energy transfer between the above-mentioned receptor and the above-mentioned G protein.

15. (currently amended) ~~Process~~ A process for detecting and/or quantifying non-covalent interactions between a target protein and one of its ligand, characterized in that:

- a fluorescent protein fused with a target protein, ~~the protein-ligand interaction of which it is desired to determine,~~ is prepared, the fusion between the fluorescent protein and said target protein being ~~[[such]]~~ so that the interaction between the target protein, ~~in particular the receptor,~~ and the ligand is not modified, and the response transduction function is not modified, the fluorescent protein being chosen from the fluorescent proteins obtained or derived from autofluorescent proteins of cnidarians, the molecular extinction coefficient of which is greater than about $14,000 \text{ M}^{-1}\text{cm}^{-1}$ and the quantic fluorescence yield of which is greater than about 0.38, this protein being further chosen from green fluorescent protein (GFP), ~~variants of GFP that conserve the fluorescence property, and fragments of GFP and of said variants that conserve the fluorescence property,~~

- said fluorescent protein fused with the target protein is placed in contact with a ligand of the above-mentioned protein, this ligand being labeled with a label comprising a molecule which is capable of absorbing the light emitted by the fluorescent protein, or a fluorescent substance, the fluorescent protein being

a fluorescence energy donor and the label being a fluorescence energy acceptor, or the fluorescent protein being a fluorescence energy acceptor and the label being a fluorescent substance which is a fluorescence energy donor, [[and]]

- irradiation is carried out at a wavelength which makes it possible either to excite the fluorescent protein or to excite the fluorescent substance, and

~~- it being possible for the above-mentioned steps of placing in contact and irradiation to be carried out either simultaneously or one after the other, or~~

~~----- said fluorescent protein fused with the target protein is placed in contact with a ligand for the above-mentioned protein, this ligand being labeled with a label comprising a molecule which is capable of absorbing the light emitted by the fluorescent protein, or a fluorescent substance, the fluorescent protein fused with the target protein or the ligand having been irradiated before being placed in contact,~~

- wherein a reduction in the amplitude of the donor's emission and or an emission signal characteristic of the acceptor's emission is detected and optionally measuring the fluorescence energy transfer when quantifying the non-covalent interactions.

16. (currently amended) A process for detecting and/or quantifying non-covalent interactions between a target protein and one of its ligands, comprising:

- preparing cells or cell fragments containing a DNA sequence comprising a [[gene]] nucleic acid expressing a fluorescent protein fused with a [[gene]] nucleic acid for the target protein, the fusion between the [[gene]] nucleic acid for the fluorescent protein and the [[gene]] nucleic acid for the target protein being [[such]] so that the properties of the target protein are not modified by the presence of the fluorescent protein, wherein the interaction between the target protein, and the ligand is not modified, and wherein a response transduction function is not modified, the fluorescent protein ~~being~~ is chosen from the fluorescent proteins obtained or derived from autofluorescent proteins of cnidarians, the molar extinction coefficient of which is greater than about $14,000 \text{ M}^{-1}\text{cm}^{-1}$ and the quantic fluorescence yield is greater than about 0.38, [[this]] said protein being selected from the group consisting of green fluorescent protein (GFP), cyan fluorescent protein (CFP), yellow fluorescent protein (YFP), and green fluorescent protein UV (GFPUV), said ligand is labeled with a label comprising a molecule capable of absorbing the light emitted by the fluorescent protein, or a fluorescent substance, the fluorescent protein being the fluorescence energy donor and the label being the fluorescence energy acceptor, or the fluorescent protein being the fluorescence

energy acceptor and the label being a fluorescent substance which is a fluorescence energy donor, and irradiating said cells or said cell fragments at a wavelength which makes it possible either to excite the fluorescent protein or to excite the fluorescent substance, wherein the steps of placing in contact and irradiating are carried out either simultaneously or one after the other, or said cells or said cell fragments are placed in contact with a ligand for said protein, said ligand labeled with a label, the cells or the ligand having been irradiated before being placed in contact,

- wherein a reduction in the amplitude of the donor's emission and/or emission signal characteristic of the acceptor's emission is detected and measuring the fluorescence energy transfer ~~when quantifying~~ to detect and/or quantify the non-covalent interactions.